

## Active Surveillance Research

---

**Educational seminars increase confidence and decreases dropout from active surveillance.** Kinsella, J. E., Van Hemelrijck, M., Allchorne, P., & Cahill, D. (2015). *European Urology Supplements*, 14(2), e363-e363a. [https://doi.org/10.1016/S1569-9056\(15\)60358-X](https://doi.org/10.1016/S1569-9056(15)60358-X)

**Factors Influencing Men's Choice of and Adherence to Active Surveillance for Low-risk Prostate Cancer: A Mixed-method Systematic Review.**

Kinsella, N., Stattin, P., Cahill, D., Brown, C., Bill-Axelson, A., Bratt, O., Carlsson, S., & Van Hemelrijck, M. (2018). *European Urology*, 74(3), 261–280. <https://doi.org/10.1016/j.eururo.2018.02.026>

**Group seminars are an effective/economic method of delivering patient information regarding radical prostatectomy and functional outcomes.**

Kinsella, J., Ashfield, A., Hazell, E., Challacombe, B., Popert, R., & Cahill, D. (2012) *International Journal of Clinical Practice*, 66.

**PCAST/SPCG-17—a randomised trial of active surveillance in prostate cancer: rationale and design.**

Ahlberg, M. S., Adami, H.-O., Beckmann, K., Bertilsson, H., Bratt, O., Cahill, D., Egevad, L., Garmo, H., Holmberg, L., & Johansson, E. (2019) *BMJ Open*, 9(8), e027860.

**Transperineal sector biopsies: Their role in active surveillance.**

Popert, R., Vyas, L., Kinsella, J., Challacombe, B., & Cahill, D. (2012). *European Urology Supplements*, 11(1), e759. [https://doi.org/10.1016/S1569-9056\(12\)60756-8](https://doi.org/10.1016/S1569-9056(12)60756-8)

**Active surveillance for prostate cancer: a systematic review of contemporary worldwide practices.**

Kinsella, N., Helleman, J., Bruinsma, S., Carlsson, S., Cahill, D., Brown, C., & Van Hemelrijck, M. (2018). *Translational Andrology and Urology*, 7(1), 83.

**As part of an active surveillance protocol transperineal saturation prostate biopsies do not compromise radical prostatectomy.**

Kinsella, J., Hemelrijck, M., Popert, R., & Cahill, D. (2012) *International Journal of Clinical Practice*, 66.

**Confirmatory biopsy for the assessment of prostate cancer in men considering active surveillance: Reference centre experience.**

Bosco, C., Kinsella, J., Acher, P., Popert, R., Cozzi, G., George, G., Cahill, D., Bianchi, R., Challacombe, B., Brown, C., Van Hemelrijck, M. *ecancermedicalscience* April 2016 DOI: 10.3332/ecancer.2016.633

**A single educational seminar increases confidence and decreases dropout from active surveillance by 5 years after diagnosis of prostate cancer.**

Kinsella, N., Beckmann, K., Cahill, D., Elhage, O., Popert, R., Cathcart, P., Challacombe, B., Brown, C., & Van Hemelrijck, M. (2019) *European Urology Oncology*, 2(4), 464–470.

**Active surveillance with confidence: the place of Transperineal Sector Biopsies (TPSB) to improve risk stratification in localised prostate cancer:**

Vyas, L., Kinsella, J., Zhao, M., Gilchrist, C., Acher, P., Chang, R., Yamamoto, H., Chandra, A., Cahill, D., & Challacombe, B. (2012) *BJU International*, 109.